Syringe Sampling Device










TKS MOUNTED ON TKC CONNECTION




TKS MOUNTED ON TKC CONNECTION

The TKS Syringe sampling device has a compact design that allows to take more samples of low viscosity products from a tank or a process line in the pharmaceutical sector with an hypodermic syringe. Once installed, the Silicone or EPDM membrane is flush positioned with the inner wall of the tank, minimizing the "Dead Leg".


Before to sampling, the port may be sterilized with alcohol and then sampling takes place by simply penetrating the special membrane with an hypodermic needle or a syringe, with minimum risk of contamination.



TKS WITH TRI-CLAMP FERRULE ASSEMBLY



TKS WITH TKL IN-LINE ASSEMBLY



TKS W/ITH TKC CONNECTION ASSEMBLY

Technical Information:

Tri-Clamp Connection: 1" 1/2 according to ASME BPE

Holes: n.10 Ø2mm

Drilled Clamp Support Material: AISI 316L - 1.4404

Blind End Cap: AISI 316L - 1.4404

Syringable gasket: White Silicone -→ complete device code TKS-AM-10X-14A

spare gasket code 40TK-X150-040

complete device code TKS-AM-10E-14A

Black EPDM spare gasket code

40TK-E150-040

Specifications:

Max pressure sample liquid: 6 barg Max sampling temperature: 95°C Not recommended for vacuum application.

Please note that the sealing capacity is reduced with increased frequency of piercing, and the membranes ability to sustain pressure decreases.It is suggested the membrane replacement every production batch.

WARNING: during the pierce, hold safely the syringe and take all the necessary precautions, considering also the pressure, temperature and the media of the sampling. DATASHE




Sampling Procedure

Sanitize the Sanitary Sampling Port immediately after installation and after each sampling operation. The sample port is designed for sampling by using an hypodermic needle as follow:

- **1**. Spray the complete assembly device with an appropriate disinfecting solution and compatible with device's materials.
- 2. Dismount the blind cap with finger-pull and place it in a disinfecting solution.
- **3**. Spray disinfecting solution on the sampling port and membrane, wait for the necessary time to obtain the desired effect.
- **4**. Look the checklist paper to identify exactly which sampling hole should be used (only one sample should be taken from each sampling hole).
- ${f 5}$. Insert the hypodermic needle for sampling. Recommended needle diameter is 0,6 1,5mm maximum.
- 6. Collect the required sample and pull out the hypodermic needle.
- **7**. Spray disinfecting solution on the sampling port and membrane.
- 8. Mount the blind cap with finger-pull.
- 9. Mark on the checklist paper which sampling hole has been used.

WARNING: during the pierce, hold safely the syringe and take all the necessary precautions, considering also the pressure, temperature and the media of the sampling.

NOTE: Pierce of membrane must be done once through the same sampling hole. Replace the used membrane when the sample port checklist is completed.



DATASHEET

